arXiv:1004.0169v2 [math.CA] 18 Jun 2010

# JACOB'S LADDERS AND THE NONLOCAL INTERACTION OF THE FUNCTION $|\zeta(1/2+it)|$ WITH THE FUNCTION $\arg\zeta(1/2+it)$ ON THE DISTANCE $\sim(1-c)\pi(t)$

JAN MOSER

Abstract. In this paper we obtain a new-type formula - *a mixed formula* - which connects the functions $|\zeta(1/2+it)|$ and $\arg\zeta(1/2+it)$. This formula cannot be obtained in the classical theory of A. Selberg, and, all the less, in the theories of Balasubramanian, Heath-Brown and Ivic.

## 1. Introduction

Let us remind that in the formula

$$\zeta\left(\frac{1}{2}+it\right)=\left|\zeta\left(\frac{1}{2}+it\right)\right|e^{i\arg\zeta\left(\frac{1}{2}+it\right)} \tag{1.1}$$

the $\arg\zeta\left(\frac{1}{2}+it\right)$ is defined as follows. If $t=\gamma$, where the $\beta+i\gamma$ is a zero of $\zeta(s)$, the $\arg\zeta\left(\frac{1}{2}+it\right)$ is obtained by continuous variation along the straight lines joining $2$, $2+it$ and $1/2+it$, starting from the value $\arg\zeta(2)=0$. If $t=\gamma$ then

$$\arg\zeta\left(\frac{1}{2}+i\gamma\right)=\lim_{t\to\gamma^+}\arg\zeta\left(\frac{1}{2}+it\right).$$

Let

$$S(t)=\frac{1}{\pi}\arg\zeta\left(\frac{1}{2}+it\right),\ S_1(T)=\int_0^T S(t)\mathrm{d}t. \tag{1.2}$$

First of all, there are the asymptotic formulae for the integrals

$$\int_0^T\left|\zeta\left(\frac{1}{2}+it\right)\right|^2\mathrm{d}t,\ \int_0^T\left|\zeta\left(\frac{1}{2}+it\right)\right|^4\mathrm{d}t,$$

(Hardy and Littlewood started to study these integrals in 1918, 1922). On the other hand, there are Selberg's asymptotic formulae ([13],[14], in 1944, 1946)

$$\int_T^{T+U}\{S(t)\}^{2k}\mathrm{d}t\sim\frac{(2k)!}{k!(2\pi)^{2k}}U(\ln\ln T)^k, \tag{1.3}$$

$$\int_T^{T+U}\{S_1(t)\}^{2k}\mathrm{d}t\sim c_kU,\ T\to\infty, \tag{1.4}$$

where $U=T^{1/2+\epsilon}$ and $k$ is a fixed positive number, (Littlewood and Titchmarsh started to study the integrals (1.3) and (1.4), $k=1$, see [2],[15] in 1925, 1928).

---

*Key words and phrases.* Riemann zeta-function.

In this paper we obtain a formula of a new type - *a mixed formula* - i.e. a formula which connects the functions (see (1.1)) $|\zeta(1/2+it)|, \arg\zeta(1/2+it)$.
This paper is a continuation of the series of papers [3]-[12].

## 2. Result

Let $\varphi_1(T),\ T\geq T_0[\varphi_1]$ stands for the Jacob's ladder. The following theorem holds true.

**Theorem**. For every fixed $k\in\mathbb{N}$ and for every fixed Jacob's ladder there is the single-valued function of $T$

$$\tau_k=\tau_k(T;\varphi_1)=\tau_k(T),\ T\geq T_1[\varphi_1],$$

for which the following asymptotic formula

$$\left|\int_0^{\varphi_1(\tau_k(T)} \arg\zeta\left(\frac{1}{2}+it\right)\mathrm{d}t\right| \sim \pi(c_k)^{\frac{1}{2k}}\frac{(\ln\tau_k(T))^{\frac{1}{2k}}}{\left|\zeta\left(\frac{1}{2}+i\tau_k(T)\right)\right|^{\frac{1}{k}}} \tag{2.1}$$

is true, where

- (A) $\tau_k\in(T,T+U),\ \varphi_1(\tau_k)\in(\varphi_1(T),\ \varphi_1(T+U)),\ U=T^{1/2+\epsilon}$
- (B) $\varphi_1(T+U)-\varphi_1(T)\sim U,\ \varphi_1(T+U)<T$
- (C) $\rho\{[T,T+U];[\varphi_1(T),\varphi_1(T+U)]\}\sim(1-c)\pi(T)\to\infty$ as $T\to\infty$ and $\rho$ denotes the distance of the corresponding segments, $c$ is the Euler's constant and $\pi(T)$ is the prime-counting function.

*Remark* 1. By (2.1) we have the prediction of the value

$$\left|\zeta\left(\frac{1}{2}+i\tau_k(T)\right)\right|=|Z(\tau_k(T))|,\ \tau_k(T)\in(T,T+U)$$

for the signal

$$Z(t)=e^{i\vartheta(t)}\zeta\left(\frac{1}{2}+it\right)$$

generated by the Riemann zeta-function, by means of the value

$$\left|\int_0^{\varphi_1(\tau_k(T))}\arg\zeta\left(\frac{1}{2}+it\right)\mathrm{d}t\right|,\ \varphi_1(\tau_k(T))\in(\varphi_1(T),\varphi_1(T+U))$$

which descends from very deep past (see (B),(C), comp. [7], Remarks 3,4).

*Remark* 2. It is quite evident that the formula (2.1) cannot be obtained within the classical theory of A. Selberg (see [13],[14]), and, all the less, in the theories of Balasubramanian, Heath-Brown and Ivic, (comp. [1]).

## 3. The first corollaries

Using the mean-value theorem in (2.1) and putting $\varphi_1(\tau_k(T))\sim\tau_k(T)$, (this follows from $t-\varphi_1(t)\sim(1-c)\pi(t)$), we obtain

**Corollary 1.**

$$|\omega[\arg\zeta]|\sim\frac{\pi(c_k)^{\frac{1}{2k}}(\ln\tau_k(T))^{\frac{1}{2k}}}{\tau_k(T)\left|\zeta\left(\frac{1}{2}+i\tau_k(T)\right)\right|^{\frac{1}{k}}}, \tag{3.1}$$

where

$$\omega[\arg\zeta]=\omega[\arg\zeta(1/2+it);\ t\in[0,\varphi_1(\tau_k(T))]]$$

denotes the mean-value of $\arg\zeta(1/2+it),\ t\in[0,\varphi_1(\tau_k(T))]$ and $\varphi_1(\tau_k(T))\in(\varphi_1(T),\varphi_1(T+U))$.

*Remark* 3. By (3.1) the following holds true: the value $|\zeta(1/2+i\tau_k(T))|$ is asymptotically defined by the mean-value $\omega[\arg\zeta]$, which descends from the very deep past (comp. Remark 1) and vice-versa.

*Remark* 4. It is evident that the set of zeroes $t=\gamma$ of the function $\zeta(1/2+it),\ t\in[T,T+U]$ is the exceptional set for the values of the function $\tau_k=\tau_k(T)$, i.e. $\tau_k(T)\neq\gamma$.

In connection with this we have the following addition

**Corollary 2.**

$$\lim_{T\to\infty}|\omega[\arg\zeta]|\cdot|\zeta(1/2+i\tau_k(T))|^{\frac{1}{k}}=0.$$

## 4. The asymptotic formula for the distance $\tilde{\gamma}'-\tilde{\gamma}$ of some subsequence of consecutive zeroes of $\zeta(1/2+it)$

4.1. For simplicity, we put $k=1$ in (3.1)

$$|\omega[\arg\zeta]|\cdot\left|\zeta\left(\frac{1}{2}+i\tau_1(T)\right)\right|\sim\pi\sqrt{c_1}\frac{\sqrt{\ln\tau_1(T)}}{\tau_1(T)}. \tag{4.1}$$

Let $t=\tilde{\gamma},\tilde{\gamma}'$ denotes the consecutive zeroes of the function $|\zeta(1/2+it)|$, for which

$$\tilde{\gamma}<\tau_1(T)<\tilde{\gamma}',\ \tau_1(T)\in(T,T+U) \tag{4.2}$$

holds true, and $n(\tilde{\gamma})$ denotes the order of the zero $t=\tilde{\gamma}$. Since

$$\begin{aligned}\left|\zeta\left(\frac{1}{2}+i\tau_1(T)\right)\right|&=|Z[\tau_1(T)]|=\\ &=\frac{1}{\{n(\tilde{\gamma})\}!}\left|Z^{(n(\tilde{\gamma}))}[\tau_1^1(T)]\right|\cdot[\tau_1(T)-\tilde{\gamma}]^{n(\tilde{\gamma})},\ \tilde{\gamma}<\tau_1^1(T)<\tau_1(T)\end{aligned} \tag{4.3}$$

(see (4.2)) then from (4.1), $(\tau_1(T)\sim\tilde{\gamma})$, the asymptotic formula

$$\tau_1(T)-\tilde{\gamma}\sim\left(\frac{\pi\sqrt{c_1}\{n(\tilde{\gamma})\}!\sqrt{\ln\tilde{\gamma}}}{\tilde{\gamma}|\omega[\arg\zeta]|\cdot|Z^{(n(\tilde{\gamma}))}[\tau_1^1(T)]|}\right)^{\frac{1}{n(\tilde{\gamma})}} \tag{4.4}$$

follows, and similarly, we obtain

$$\begin{aligned}&|\tau_1(T)-\tilde{\gamma}'|=\tilde{\gamma}'-\tau_1(T)\sim\\ &\left(\frac{\pi\sqrt{c_1}\{n(\tilde{\gamma}')\}!\sqrt{\ln\tilde{\gamma}'}}{\tilde{\gamma}'|\omega[\arg\zeta]|\cdot|Z^{(n(\tilde{\gamma}'))}[\tau_1^1(T)]|}\right)^{\frac{1}{n(\tilde{\gamma}')}},\ \tau_1(T)<\tau_1^2(T)<\tilde{\gamma}'.\end{aligned} \tag{4.5}$$

Then we obtain from (4.4), (4.5)

**Corollary 3.** If the pair of consecutive zeroes $\tilde{\gamma},\tilde{\gamma}'$ fulfils (4.2) then the asymptotic formula

$$\begin{aligned}\tilde{\gamma}'-\tilde{\gamma}\sim&\left(\frac{\pi\sqrt{c_1}\{n(\tilde{\gamma})\}!\sqrt{\ln\tilde{\gamma}}}{\tilde{\gamma}|\omega[\arg\zeta]|\cdot|Z^{(n(\tilde{\gamma}))}[\tau_1^1(T)]|}\right)^{\frac{1}{n(\tilde{\gamma})}}+\\ &+\left(\frac{\pi\sqrt{c_1}\{n(\tilde{\gamma}')\}!\sqrt{\ln\tilde{\gamma}'}}{\tilde{\gamma}'|\omega[\arg\zeta]|\cdot|Z^{(n(\tilde{\gamma}'))}[\tau_1^1(T)]|}\right)^{\frac{1}{n(\tilde{\gamma}')}},\ \tilde{\gamma}<\tau_1^1(T)<\tau_1(T)<\tau_1^2(T)<\tilde{\gamma}'\end{aligned} \tag{4.6}$$

holds true.

*Remark* 5. This is the surprise (for the author) that the asymptotic formula for $\tilde{\gamma}'-\tilde{\gamma}$ contains the expression $|\omega[\arg\zeta]|$.

*Remark* 6. The nonlocal nature of the formula (4.6) is quite evident (comp. Remark 1).

4.2. Let $\{(\bar{\gamma},\bar{\gamma}')\}$ denotes the subsequence of the sequence $\{(\tilde{\gamma},\tilde{\gamma}')\}$ such that $n(\bar{\gamma})=n(\bar{\gamma}')=1$ (for example if the weakened Mertens hypothesis is true). Then from (4.6) we obtain

**Corollary 4.**

$$\bar{\gamma}'-\bar{\gamma}\sim\frac{\pi\sqrt{c_1}\sqrt{\ln\bar{\gamma}}}{\bar{\gamma}|\omega[\arg\zeta]|}\left(\frac{1}{|Z'[\tau_1^1]|}+\frac{1}{|Z'[\tau_1^2]|}\right),\ \bar{\gamma}<\tau_1^1<\tau_1<\tau_1^2<\bar{\gamma}'. \tag{4.7}$$

Next, using the estimate

$$|Z'|<A\bar{\gamma}^{1/6}\ln\bar{\gamma},\ \bar{\gamma}'-\bar{\gamma}<A\bar{\gamma}^{1/6+\epsilon/2} \tag{4.8}$$

we obtain from (4.7) the following

**Corollary 5.**

$$|\omega[\arg\zeta]|>\frac{A}{\bar{\gamma}^{4/3+\epsilon}},\ \bar{\gamma}\to\infty. \tag{4.9}$$

*Remark* 7. The estimate (4.9) is the first nontrivial lower bound of $|\omega[\arg\zeta]|$, (the small improvements of the exponent 1/6 in (4.8) are irrelevant).

## 5. Lemmas

5.1. Let us remind that

$$\tilde{Z}^2(t)=\frac{\mathrm{d}\varphi_1(t)}{\mathrm{d}t},\ \varphi_1(t)=\frac{1}{2}\varphi(t), \tag{5.1}$$

where

$$\tilde{Z}^2(t)=\frac{Z^2(t)}{2\Phi'_\varphi[\varphi(t)]}=\frac{Z^2(t)}{\left\{1+\mathcal{O}\left(\frac{\ln\ln t}{\ln t}\right)\right\}\ln t}, \tag{5.2}$$

(see [3], (3.9); [5], (1.3); [9], (1.1), (3.1), (3.2)). The following Lemma holds true (see [8], (2.5); [9], (3.3)).

**Lemma 1.** *For every integrable function (in the Lebesgue sense) $f(x)$, $x\in[\varphi_1(T),\varphi_1(T+U)]$ the following is true*

$$\int_T^{T+U}f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{\varphi_1(T)}^{\varphi_1(T+U)}f(x)\mathrm{d}x,\ U\in\left(0,\frac{T}{\ln T}\right], \tag{5.3}$$

*where $t-\varphi_1(t)\sim(1-c)\pi(t)$.*

*Remark* 8. The formula (5.3) remains true also in the case when the integral on the right-hand side of (5.3) is only relatively convergent improper integral of the second kind (in the Riemann sense).

5.2. Next, the following $\tilde{Z}^2$-transformation of the formulae of A. Selberg (1.3), (1.4) holds true (comp. [7], Concluding remarks).

**Lemma 2.**

$$\int_T^{T+U}\{S[\varphi_1(t)]\}^{2k}\left|\zeta\left(\frac{1}{2}+it\right)\right|^2\mathrm{d}t\sim\frac{(2k)!}{k!(2\pi)^{2k}}U\ln T(\ln\ln T)^k, \tag{5.4}$$

$$\int_T^{T+U}\{S_1[\varphi_1(t)]\}^{2k}\left|\zeta\left(\frac{1}{2}+it\right)\right|^2\mathrm{d}t\sim c_kU\ln T. \tag{5.5}$$

*Proof.* From (1.3) and by Lemma 1 we obtain

$$\begin{aligned}\int_T^{T+U}\{S[\varphi_1(t)]\}^{2k}\tilde{Z}^2(t)\mathrm{d}t&=\int_{\varphi_1(T)}^{\varphi_1(T+U)}\{S[\varphi_1(T)]\}^{2k}\mathrm{d}t\sim\\ &\sim\frac{(2k)!}{k!(2\pi)^{2k}}\{\varphi_1(T+U)-\varphi_1(T)\}\ln\ln\varphi_1(T)=\\ &=\frac{(2k)!}{k!(2\pi)^{2k}}U\frac{\varphi_1(T+U)-\varphi_1(T)}{U}\ln\ln\varphi_1(T)=\\ &=\frac{(2k)!}{k!(2\pi)^{2k}}U\tan[\alpha(T,U)]\ln\ln\varphi_1(T),\end{aligned} \tag{5.6}$$

where $\alpha(T,U)$ is the angle of the chord of the curve $y=\varphi_1(t)$ that binds the points $[T,\varphi_1(T)]$ and $[T+U,\varphi_1(T+U)]$. Let us remind that

$$\tan[\alpha(T,U)]\sim 1,\ U\in\left[T^{1/3+\epsilon},\frac{T}{\ln T}\right], \tag{5.7}$$

(comp. [3], (8.3), [7], (3.9)) and

$$t-\varphi_1(t)\sim(1-c)\pi(t), \tag{5.8}$$

(see (5.1), [3], (6.2)). Since by (5.8) we have

$$\varphi_1(T)\sim T,\ \ln\ln\varphi_1(T)\sim\ln\ln T, \tag{5.9}$$

then from (5.6) by (5.7), (5.9) this formula

$$\int_T^{T+U}\{S[\varphi_1(t)]\}^{2k}\tilde{Z}^2(t)\mathrm{d}t\sim\frac{(2k)!}{k!(2\pi)^{2k}}U(\ln\ln T)^k \tag{5.10}$$

follows. Using the mean-value theorem in (5.10) we obtain (5.4) by (5.2). Similarly we obtain (5.5). □

## 6. Proof of the Theorem

Using the mean-value theorem in (5.5) we obtain

$$\begin{aligned}&|S_1[\varphi_1(\tau)]|^{2k}Z^2(\tau)\sim c_k\ln T\sim c_k\ln\tau,\\ &\tau=\tau(T,U,k)\in(T,T+U),\ \varphi_1(\tau)\in(\varphi_1(T),\varphi_1(T+U)).\end{aligned} \tag{6.1}$$

Since $U=T^{1/2+\epsilon}$ then $\tau=\tau(T,k,\epsilon),\ T\geq T_0[\varphi_1]$. Next, if for every $T\geq T_0[\varphi_1]$ we take one mean-value $\tau_k(T)\in\{\tau(T,k,\epsilon)\}$ then we have the single-valued function $\tau_k=\tau_k(T)$ for fixed $k,\epsilon$ (comp. [7], Remark 2). Hence by (5.1) we have

$$\begin{aligned}&|S_1[\varphi_1(\tau_k(T))]|^{2k}\sim c_k\frac{\ln\tau_k(T)}{\left|\zeta\left(\frac{1}{2}+i\tau_k(T)\right)\right|^2},\\ &\tau_k(T)\in(T,T+U),\ \varphi_1(\tau_k(T))\in(\varphi_1(T),\varphi_1(T+U)).\end{aligned} \tag{6.2}$$

Then from (6.2) the asymptotic formula (2.1) and (A) follows. The expressions (B) and (C) are identical with (C), (D), $k = 1$ of Theorem in [9].

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

## REFERENCES

[1] A. Ivic, ‘The Riemann zeta-function‘, A Willey-Interscience Publication, New York, 1985.

[2] J.E. Littlewood, ‘On the Riemann zeta-function’, Proc. Lond. Math. Soc. (2), 24, (1925), 175-201.

[3] J. Moser, ‘Jacob's ladders and the almost exact asymptotic representation of the Hardy-Littlewood integral’, (2008), arXiv:0901.3973.

[4] J. Moser, ‘Jacob's ladders and the tangent law for short parts of the Hardy-Littlewood integral’, (2009), arXiv:0906.0659.

[5] J. Moser, ‘Jacob's ladders and the multiplicative asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral’, (2009), arXiv:0907.0301.

[6] J. Moser, ‘Jacob's ladders and the quantization of the Hardy-Littlewood integral’, (2009), arXiv:0909.3928.

[7] J. Moser, ‘Jacob's ladders and the first asymptotic formula for the expression of the sixth order $|\zeta(1/2+i\varphi(t)/2)|^4|\zeta(1/2+it)|^2$’, (2009), arXiv:0911.1246.

[8] J. Moser, ‘Jacob's ladders and the first asymptotic formula for the expression of the fifth order $Z[\varphi(t)/2+\rho_1]Z[\varphi(t)/2+\rho_2]Z[\varphi(t)/2+\rho_3]\hat{Z}^2(t)$ for the collection of disconnected sets‘, (2009), arXiv:0912.0130.

[9] J. Moser, ‘Jacob's ladders, the iterations of Jacob's ladder $\varphi_1^k(t)$ and asymptotic formulae for the integrals of the products $Z^2[\varphi_1^n(t)]Z^2[\varphi^{n-1}(t)]\cdots Z^2[\varphi_1^0(t)]$ for arbitrary fixed $n \in \mathbb{N}$‘ (2010), arXiv:1001.1632.

[10] J. Moser, ‘Jacob's ladders and the asymptotic formula for the integral of the eight order expression $|\zeta(1/2+i\varphi_2(t))|^4|\zeta(1/2+it)|^4$‘, (2010), arXiv:1001.2114.

[11] J. Moser, ‘Jacob's ladders and the asymptotically approximate solutions of a nonlinear diophantine equation‘, (2010), arXiv: 1001.3019.

[12] J. Moser, ‘Jacob's ladders and the asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral of the function $|\zeta(1/2+it)|^4$‘, (2010), arXiv:1001.4007.

[13] A. Selberg, ‘On the remainder in the formula for $N(T)$, the number of zeros of $\zeta(s)$ in the strip $0 < t < T$‘, Avh. Norsk. Vid. Akad. Oslo (1944), No.1. 1-27.

[14] A. Selberg, ‘Contributions to the theory of the Riemann zeta-function‘, Arch. for Math. ogNaturv. B, 48 (1946), No. 5, 89-155.

[15] E.C. Titchmarsh, ‘On the remainder in the formula for $N(T)$ the number of zeros in the strip $0 < t < T$‘, Proc. London Math. Soc. (2), 27 (1928), 449-458.

DEPARTMENT OF MATHEMATICAL ANALYSIS AND NUMERICAL MATHEMATICS, COMENIUS UNIVERSITY, MLYNSKA DOLINA M105, 842 48 BRATISLAVA, SLOVAKIA
*E-mail address*: `jan.mozer@fmph.uniba.sk`